Table 1. Global Prospective HoLISTIC Hodgkin Lymphoma Clinical Trials (individual patient data).

Group/Study Name (ref)	Years accrual	Pt number	Author, Year(s) published	Reference(s)	
Adult Early Stage					
EORTC H9U	1998 – 2002	808	Ferme 2017	[1]	
NCI-C/ECOG	1994 – 2002	276	Meyer 2005 and Meyer 2012	[2, 3]	
*ECOG 2496	1999 – 2006	281	Gordon 2012, Evens 2013, and Advani 2015	[4-6]	
EORTC H10	2006 – 2009	1950	Raemaekers 2014 and Andre 2017	[7, 8]	
RAPID	2003 – 2010	602	Radford 2015	[9]	
*RATHL	2008 – 2012	500	Johnson 2016	[10]	
Pediatric Low and Interm	ediate Risk (stages I-II)				
AHOD0031	2002 – 2009	1114	Friedman 2014	[11]	
AHOD0431	2006 – 2012	287	Keller 2018	[12]	
Adult Early Stage and Pediatric Low-Intermediate Risk (total): 5,818 patients					
Adult Advanced Stage					
Italian IIL HD9601	1996 – 2000	355	Gobbi 2005 and Chisesi 2011	[13, 14]	
Italian HD2000	2000 – 2007	307	Federico 2009 and Merli 2016	[15, 16]	
UK ISRCTN	1998 – 2006	520	Hoskin 2009	[17]	
*ECOG 2496	1999 – 2006	513	Gordon 2012, Evens 2013, and Advani 2015	[4-6]	
SWOG 0816	2009 – 2012	371	Press 2016 and Stephens 2019	[18, 19]	
*RATHL	2008 – 2012	712	Johnson 2016	[10]	
HD0801 IIL	2008 – 2013	519	Zinzani 2017	[20]	
GITIL HD0607	2008 – 2014	780	Gallamini 2018	[21]	
Pediatric Intermediate an	d High Risk (stages III-IV)	)			
AHOD0031	2002 - 2009	608	Friedman 2014	[11]	
AHOD0831	2009 – 2013	165	Kelly 2019	[22]	
Adult Advanced Stage an	nd Pediatric Intermediate-	High Risk (total	): 4,850 patients		
*adult studies that included					

Table 2. Global HoLISTIC Hodgkin Lymphoma Registries and Prospective Cohorts (individual patient data).

Registry Name (ref)	No. Pts	Yrs Diagnosis	Author, Year(s) published	Reference
Princess Margaret Hospital	1200	2000-2017	Hahn 2017	[23]
Iowa/Mayo Clinic SPORE	700	2002-2017	Cerhan 2017	[24]
BC Cancer	1905	2000-2015	Hapgood 2016	[25]
Australia	470	2016-present	Lymphoma, Related Diseases Registry 2022	[26]
Stanford Registry*	395	1995-2019	n/a	n/a
TOTAL: 4275 pto				

**TOTAL: 4275 pts** 

Abbreviations: Ref, references; No, number; pts; patients; BC, British Columbia.

\*Early stage only; transfer of advanced stage patients in process

## References

- Ferme, C., et al., ABVD or BEACOPPbaseline along with involved-field radiotherapy in early-stage Hodgkin Lymphoma with risk factors: Results of the European Organisation for Research and Treatment of Cancer (EORTC)-Groupe d'Etude des Lymphomes de l'Adulte (GELA) H9-U intergroup randomised trial. Eur J Cancer, 2017. 81: p. 45-55.
- 2. Meyer, R.M., et al., Randomized comparison of ABVD chemotherapy with a strategy that includes radiation therapy in patients with limited-stage Hodgkin's lymphoma: National Cancer Institute of Canada Clinical Trials Group and the Eastern Cooperative Oncology Group. J Clin Oncol, 2005. **23**(21): p.4634-42.
- 3. Meyer, R.M., et al., *ABVD alone versus radiation-based therapy in limited-stage Hodgkin's lymphoma*. N Engl J Med, 2012. **366**(5): p. 399-408.
- 4. Gordon, L.I., et al., Randomized phase III trial of ABVD versus Stanford V with or without radiation therapy in locally extensive and advanced-stage Hodgkin lymphoma: an intergroup study coordinated by the Eastern Cooperative Oncology Group (E2496). J Clin Oncol, 2013. **31**(6): p. 684-91.
- 5. Advani, R.H., et al., Randomized Phase III Trial Comparing ABVD Plus Radiotherapy With the Stanford V Regimen in Patients With Stages I or II Locally Extensive, Bulky Mediastinal Hodgkin Lymphoma: A Subset Analysis of the North American Intergroup E2496 Trial. J Clin Oncol, 2015. 33(17): p. 1936-42.
- 6. Evens, A.M., et al., *The efficacy and tolerability of adriamycin, bleomycin, vinblastine, dacarbazine and Stanford V in older Hodgkin lymphoma patients: a comprehensive analysis from the North American intergroup trial E2496.* Br J Haematol, 2013. **161**(1): p. 76-86.
- 7. Andre, M.P.E., et al., *Early Positron Emission Tomography Response-Adapted Treatment in Stage I and II Hodgkin Lymphoma: Final Results of the Randomized EORTC/LYSA/FIL H10 Trial.* J Clin Oncol, 2017. **35**(16): p. 1786-1794.
- 8. Raemaekers, J.M., et al., Omitting radiotherapy in early positron emission tomography-negative stage I/II Hodgkin lymphoma is associated with an increased risk of early relapse: Clinical results of the preplanned interim analysis of the randomized EORTC/LYSA/FIL H10 trial. J Clin Oncol, 2014. **32**(12): p. 1188-94.
- 9. Radford, J., et al., Results of a trial of PET-directed therapy for early-stage Hodgkin's lymphoma. N. Engl. J. Med, 2015. **372**(17): p. 1598-1607.
- Johnson, P., et al., Adapted Treatment Guided by Interim PET-CT Scan in Advanced Hodgkin's Lymphoma.
  N Engl J Med, 2016. 374(25): p. 2419-29.
- 11. Keller, F.G., et al., Results of the AHOD0431 trial of response adapted therapy and a salvage strategy for limited stage, classical Hodgkin lymphoma: A report from the Children's Oncology Group. Cancer, 2018.
- 12. Friedman, D.L., et al., Dose-intensive response-based chemotherapy and radiation therapy for children and adolescents with newly diagnosed intermediate-risk hodgkin lymphoma: a report from the Children's Oncology Group Study AHOD0031. J Clin Oncol, 2014. **32**(32): p. 3651-8.
- 13. Chisesi, T., et al., Long-term follow-up analysis of HD9601 trial comparing ABVD versus Stanford V versus MOPP/EBV/CAD in patients with newly diagnosed advanced-stage Hodgkin's lymphoma: a study from the Intergruppo Italiano Linfomi. J Clin Oncol, 2011. **29**(32): p. 4227-33.
- 14. Gobbi, P.G., et al., ABVD versus modified stanford V versus MOPPEBVCAD with optional and limited radiotherapy in intermediate- and advanced-stage Hodgkin's lymphoma: final results of a multicenter randomized trial by the Intergruppo Italiano Linfomi. J Clin Oncol, 2005. **23**(36): p.9198-207.
- 15. Federico, M., et al., ABVD compared with BEACOPP compared with CEC for the initial treatment of patients with advanced Hodgkin's lymphoma: results from the HD2000 Gruppo Italiano per lo Studio dei Linfomi Trial. J Clin Oncol, 2009. **27**(5): p. 805-11.
- Merli, F., et al., Long-Term Results of the HD2000 Trial Comparing ABVD Versus BEACOPP Versus COPP-EBV-CAD in Untreated Patients With Advanced Hodgkin Lymphoma: A Study by Fondazione Italiana Linfomi. J Clin Oncol, 2016. 34(11): p. 1175-81.
- 17. Hoskin, P.J., et al., Randomized comparison of the stanford V regimen and ABVD in the treatment of advanced Hodgkin's Lymphoma: United Kingdom National Cancer Research Institute Lymphoma Group Study ISRCTN 64141244. J Clin Oncol, 2009. **27**(32): p. 5390-6.
- 18. Press, O.W., et al., US Intergroup Trial of Response-Adapted Therapy for Stage III to IV Hodgkin Lymphoma Using Early Interim Fluorodeoxyglucose-Positron Emission Tomography Imaging: Southwest Oncology Group S0816. J Clin Oncol, 2016. **34**(17): p. 2020-7.

- 19. Stephens, D.M., et al., Five-year follow-up of SWOG S0816: limitations and values of a PET-adapted approach with stage III/IV Hodgkin lymphoma. Blood, 2019. **134**(15): p. 1238-1246.
- Zinzani, P.L., et al., Interim Positron Emission Tomography Response-Adapted Therapy in Advanced-Stage Hodgkin Lymphoma: Final Results of the Phase II Part of the HD0801 Study. J Clin Oncol, 2016. 34(12): p. 1376-85.
- 21. Gallamini, A., et al., Early Chemotherapy Intensification With Escalated BEACOPP in Patients With Advanced-Stage Hodgkin Lymphoma With a Positive Interim Positron Emission Tomography/Computed Tomography Scan After Two ABVD Cycles: Long-Term Results of the GITIL/FIL HD 0607 Trial. J Clin Oncol, 2018. 36(5): p. 454-462.
- 22. Kelly, K.M., et al., Response-adapted therapy for the treatment of children with newly diagnosed high risk Hodgkin lymphoma (AHOD0831): a report from the Children's Oncology Group. Br J Haematol, 2019. **187**(1): p. 39-48.
- 23. Hahn, E., et al., Late Cardiac Toxicity After Mediastinal Radiation Therapy for Hodgkin Lymphoma: Contributions of Coronary Artery and Whole Heart Dose-Volume Variables to Risk Prediction. Int J Radiat Oncol Biol Phys, 2017. **98**(5): p. 1116-1123.
- 24. Cerhan, J.R., et al., Cohort Profile: The Lymphoma Specialized Program of Research Excellence (SPORE) Molecular Epidemiology Resource (MER) Cohort Study. Int J Epidemiol, 2017. **46**(6): p. 1753-1754i.
- 25. Hapgood, G., et al., Evaluation of the Risk of Relapse in Classical Hodgkin Lymphoma at Event-Free Survival Time Points and Survival Comparison With the General Population in British Columbia. J Clin Oncol, 2016. **34**(21): p. 2493-500.
- 26. Lymphoma and I. Related Diseases Registry, *Improving outcomes for patients with lymphoma: design and development of the Australian and New Zealand Lymphoma and Related Diseases Registry.* BMC Med Res Methodol, 2022. **22**(1): p. 266.